



























42. Carmont MR, Maffulli N. Medial patellofemoral ligament reconstruction: a new technique. *BMC Musculoskelet Disord*. 2007;8:22.
43. Matsushita T, Kuroda R, Araki D, et al. Medial patellofemoral ligament reconstruction with lateral soft tissue release in adult patients with habitual patellar dislocation. *Knee Surg Sports Traumatol Arthrosc*. 2013;21(3):726-30.
44. Sandmeier RH, Burks RT, Bachus KN, et al. The effect of reconstruction of the medial patellofemoral ligament on patellar tracking. *Am J Sports Med*. 2000;28: 345-9.
45. Baker RH, Carroll N, Dewar FP, et al. The semitendinosus tenodesis for recurrent dislocation of the patella. *J Bone Joint Surg Br*. 1972;54(1):103-9.
46. Grannatt K, Heyworth BE, Ogunwole O, Micheli LJ, Kocher MS. Galeazzi semitendinosus tenodesis for patellofemoral instability in skeletally immature patients. *J Pediatr Orthop*. 2012;32(6):621-5.
47. Pradhan RL, Watanabe W, Itoi E, et al. Total knee arthroplasty in bilateral congenital dislocation of the patella-a case report. *Acta Orthop Scand*. 2001;72(4):422-4.
48. Bauer J, Martus J. Pediatric Orthopaedic Society of North America. Congenital Dislocation of the Patella Study Guide. Available at: <https://posna.org/Physician-Education/Study-Guide/Congenital-Dislocation-of-Patella>. Accessed April 1, 2020.
49. Chang CB, Shetty GM, Lee JS, et al. A Combined Closing Wedge Distal Femoral Osteotomy and Medial Reefing Procedure for Recurrent Patellar Dislocation with Genu Valgum. *Yonsei Med J*. 2017;58(4):878–883.
50. Bullek DD, Scuderi GR, Insall JN. Management of the chronic irreducible patellar dislocation in total knee arthroplasty. *J Arthroplasty*. 1996 11(3):339–345
51. Drexler M, Dwyer T, Dolkart O, et al. Tibial rotational osteotomy and distal tuberosity transfer for patella subluxation secondary to excessive external tibial torsion: surgical technique and clinical outcome. *Knee Surg Sports Traumatol Arthrosc*. 2014;22(11):2682-9.
52. Ramaswamy R, Kosashvili Y, Murnaghan JJ, et al. Bilateral rotational osteotomies of the proximal tibiae and tibial tuberosity distal transfers for the treatment of congenital lateral dislocations of patellae: a case report and literature review. *Knee*. 2009;16(6):507-11.